

Ecuador

Medical Summary

General Information

Ecuador is a developing nation but is in the upper half of the world's economies. Located along the equator on the west coast of South America, its climate is tropical along the coast and in jungle lowlands, and cooler inland at higher elevations.

Travel Immunizations

Hepatitis A

Recommended for: all travelers.

Typhoid fever

Recommended for: most travelers, especially those with adventurous dietary habits; those without consistent access to safe food and water; those with prolonged stays; and those traveling outside common tourist packages and other prearranged fixed itineraries, especially in rural areas.

Consider for: all risk-averse travelers desiring maximum pretravel preparation.

Influenza

Risk exists from June through September and from January through March, although off-season transmission can occur.

Recommended for: all travelers during transmission season due to demonstrated influenza risk in this group.

Travelers not already immunized with the currently available vaccine formulation should be vaccinated. Travelers immunized with the current formulation more than 6 months earlier should consider revaccination because immunity may have declined. Consider oseltamivir as standby therapy, especially for those who are at high risk for complications from influenza or inadequately vaccinated.

Hepatitis B

Recommended for: prolonged stays; frequent short stays in this or other high-risk countries; adventure travelers; the possibility of acupuncture, dental work, or tattooing; all health care workers; the possibility of a new sexual partner during the stay; injection drug users; and travelers with high potential to seek medical care in local facilities.

Increased awareness is recommended regarding safe sex and body fluid/blood precautions.

Yellow fever

Requirement (for entry, per WHO): A vaccination certificate is required for travelers aged ≥ 1 year coming from Brazil, Democratic Republic of the Congo, and Uganda. Note: This applies to airport layovers longer than 12 hours in these countries.

Recommendation (for health protection):

Recommended for travelers aged ≥ 9 months: areas east of the Andes Mountains below 2,300 m (7,500 ft) (see map). Daytime insect precautions are essential for unvaccinated travelers.

Not recommended except for highly risk-averse travelers and long-stay travelers: itineraries limited to areas west of the Andes Mountains below 2,300 m except Guayaquil (see map). No human cases of YF have ever been reported from these areas, and data analysis by WHO indicates extremely low potential for YF virus exposure.

Not recommended: itineraries limited to the cities of Guayaquil, Quito, Cuenca, or Otavalo; the Cotopaxi Volcano; areas above 2,300 m; or the Galápagos Islands.

Measles, mumps, rubella

Indicated for those born in 1957 or later (1970 or later in Canada and U.K.; 1966 or later in Australia) without evidence of immunity or of 2 countable doses of live vaccine at any time during their lives. Also indicated for those born before 1970 (in

Canada) without evidence of immunity or previous vaccination with 1 countable dose of measles-containing vaccine.

Rabies

Significant risk from bats exists in most jungle areas. Canine rabies rarely occurs.

Recommended for prolonged stays: all travelers and expatriates, with a priority for young children.

Recommended for short stays: adventure travelers, hikers, cyclists, and cavers; locations more than 24 hours' travel from a reliable source of human rabies immune globulin and rabies vaccine for postexposure treatment; animal workers (such as veterinarians and wildlife professionals); and all travelers likely to have contact with bats.

Consider for: risk-averse travelers with short stays desiring maximum pretravel preparation.

Dog, other terrestrial mammal, and bat bites or scratches should be taken seriously, and postexposure prophylaxis should be sought even in those already immunized.

The Galapagos Islands only: Risk from bats is presumed to exist throughout the island. Rabies is not present in canines or other mammals.

Recommended for: all travelers likely to have contact with bats.

Bat bites or scratches should be taken seriously, and postexposure prophylaxis should be sought even in those already immunized.

Routine Immunizations

Tetanus, diphtheria, pertussis

Due to increasingly frequent pertussis outbreaks worldwide, all travelers should receive Tdap vaccine every 10 years, assuming they previously received an adequate primary series. Those who received Td or TT for their most recent booster should receive an immediate dose of Tdap, regardless of the interval since the last tetanus dose.

Pneumococcal

Recommended for adults aged ≥ 65 years and all adults with chronic disease or immunocompromising conditions.

Varicella

Indicated for all persons born outside the U.S. or born in the U.S. after 1979, except not indicated for persons with an adequate vaccination history (2 lifetime doses), reliable evidence of previous infection, or laboratory confirmation of immunity.

Malaria

General malaria information: predominantly *P. vivax*. Transmission occurs throughout the year.

Location-specific recommendations:

Chemoprophylaxis is recommended for all travelers: altitudes below 1,500 m (4,900 ft) in Carchi Province and in certain cantons of Esmeraldas, Sucumbios, Orellana, Pastaza, Morona-Santiago, and Azuay provinces; all cities and towns within these areas except the central urban area of Esmeraldas.

Chemoprophylaxis is recommended for certain travelers (see Issues to Consider box): altitudes below 1,500 m in certain cantons of Esmeraldas, Napo, Cotopaxi, Bolívar, Los Ríos, Guayas, Sucumbios, Orellana, and Morona-Santiago provinces; all cities and towns within these areas except the central urban area of Tena.

Insect precautions only are recommended (negligible transmission is reported): the central urban areas of Esmeraldas and Tena; altitudes below 1,500 m in various provinces throughout the country; all cities and towns within these areas except Guayaquil, Manta, Portoviejo, and Babahoyo.

No preventive measures are necessary (no evidence of transmission exists): Quito, Manta, Babahoyo, Guayaquil, Portoviejo, and all other cities and towns not mentioned above; the Galápagos Islands; the provinces of Chimborazo, Santa Elena, and Loja; altitudes above 1,500 m; all other areas not mentioned above.

Preventive measures: Evening and nighttime insect precautions are essential in areas with any level of transmission. Atovaquone-proguanil (Malarone or generic), doxycycline, and mefloquine are protective in this country. Drug choice depends on personal factors discussed between the traveler and medical provider.

No preventive measure is 100% effective. Immediate medical attention is necessary for fever or influenza-like illness within 3 months after travel in a malaria risk area. Include mention of travel history.

Issues to Consider	
<i>Factors favoring chemoprophylaxis</i>	<i>Factors against chemoprophylaxis</i>
<ul style="list-style-type: none"> Adventure travel Risk-averse and vulnerable travelers Areas subject to infrequent epidemics Immigrants visiting friends and relatives Flexible itineraries Travel longer than 1 month Unreliable medical expertise and/or treatment drugs at destination 	<ul style="list-style-type: none"> Air-conditioned hotels only Urban areas only Non-transmission season Minimal nighttime exposure Travel shorter than 3 days
<p>For more information, see <i>Technical Explanation of Malaria Mapping</i>.</p>	

Travelers' Diarrhea

High risk exists throughout the country, with moderate risk in deluxe accommodations. Food and beverage precautions may reduce the likelihood of illness.

Travelers should carry loperamide for self-treatment of diarrhea and azithromycin to add if diarrhea is severe.

Other Concerns

Altitude illness

Chemoprophylaxis with acetazolamide should be considered for travelers anticipating rapid ascent to sleeping altitudes above 2,800 m (9,200 ft). The elevation of Quito is 2,800 m. Climbers summiting Cotopaxi will reach an elevation of 6,200 m (20,300 ft).

Dengue

Risk exists in urban and rural areas throughout the country at elevations below 2,300 m (7,500 ft), including on the Galápagos Islands, especially in Manabí and Guayas provinces. No risk exists in Quito. Transmission occurs throughout the year, especially during the rainy season, with highest activity from January through September. Daytime insect precautions are recommended.

Chikungunya

Low risk exists throughout most of the country at elevations below 2,300 m (7,500 ft), mainly in Guayas Province. Transmission occurs throughout the year. Daytime insect precautions are recommended.

Zika

Risk exists in Sucumbíos Province and areas west of the Andes Mountains at elevations below 2,300 m (7,500 ft), especially in Manabí Province. Low risk exists on the Galápagos Islands. No risk exists in Quito. Daytime insect precautions are recommended. Pregnant women (in any trimester) from nonaffected areas should not travel to affected areas with evidence of virus circulation after 2015.

Marine hazards

Risk from jellyfish exists, including highly venomous bluebottle jellyfish. Travelers wading, launching boats, or fishing are especially at risk. Risk from stonefish and sea urchins exists. Risk from coral is limited to Galápagos Islands. Travelers should seek out and heed posted warnings and refrain from bathing at unmarked, unpatrolled beaches.

Tuberculosis

Tuberculosis (TB) is common in all developing countries and presents risk in certain developed countries. TB incidence in this country is 25 to 100 cases per 100,000 population (not the highest risk category).

A documented interferon gamma release assay (IGRA) or a tuberculin skin test (TST) is recommended before departure and after return for stays longer than 1 month for health care workers and those with anticipated exposure in prisons, homeless shelters, refugee camps, or shanty towns.

Travelers should avoid public transportation and people who are coughing in crowded, public places (whenever possible). Domestic help should be screened for TB.

Leishmaniasis

Risk of cutaneous and mucosal disease exists throughout the country at elevations below 3,000 m (9,800 ft), especially in northern and Amazonian provinces. No risk exists on the Galápagos Islands. Evening and nighttime sandfly precautions are recommended.

Air pollution

Air quality may be variable throughout the year. Annual mean particulate matter concentrations are unhealthy.

Santo Domingo de Los Colorados: When air quality worsens, travelers should reduce prolonged or heavy outdoor exertion; those with lung disease or at the extremes of age should avoid prolonged or heavy outdoor exertion.

Latacunga or Quito: When air quality worsens, travelers with lung disease or at the extremes of age should reduce prolonged or heavy outdoor exertion.

Snakebites

Risk of envenomation exists in areas with dense vegetation or rock formations (especially in warm weather when snakes tend to be more active). Most snakebites result from startling snakes; do not disturb or handle snakes. Boots and long pants are recommended in high-threat situations. Medical care is indicated after any snakebite.

Chagas' disease (American trypanosomiasis)

Risk to travelers is unknown but is presumed to be low in rural areas throughout the country. Travelers should avoid overnight stays in houses constructed of mud, adobe brick, or palm thatch.

Leptospirosis

Risk exists and is presumed to have widespread distribution. Travelers who anticipate activities with extensive outdoor exposure (e.g., hiking, biking, swimming, or rafting) should consider weekly prophylaxis with doxycycline.

Plague

Negligible risk exists and is limited to Chimborazo, Cotopaxi, and Loja provinces. Transmission occurs throughout the year. Travelers should avoid contact with potentially infected rodents and their fleas.

Bartonellosis

Risk exists mainly in Loja, Guayas, and Zamora-Chinchipec provinces and less commonly in Manabí Province. Risk to travelers is low. Insect precautions are recommended.

Security

Consular Travel Warning: Due to drug trafficking and other ongoing security concerns, Canada (GAC) advises avoiding travel to areas bordering Colombia and Peru and advises reconsidering travel (or avoiding nonessential travel) to Montañita (Santa Elena Province). U.S. (DOS), U.K. (FCO), and Australia (DFAT) have more limited warnings.

Key Security Threats: Violent crime, petty crime, and express kidnappings occur throughout the country (especially in Quito, Guayaquil, and other cities) and on volcano hiking trails (including Cerro Mandango near Vilcabamba, Loja Province). Kidnappings occur in northern and northeastern areas bordering Colombia and Peru, including the Cuyabeno Wildlife Reserve. Foreigners, including those working for oil companies, may be targeted. Additional threats to personal safety include unsafe passenger boats in the Galápagos Islands. For more information, see Consular Advice.

Aviation Assessment: U.S. Federal Aviation Administration has determined that the civil aviation authority of this country oversees its air carriers in accordance with minimum international safety standards.

Medical Care

Adequate private medical care that meets most international standards is available in Quito, Cuenca and Guayaquil. Medical care throughout the rest of the country is inadequate and usually does not meet international standards. One or more JCI accredited hospitals are present in this country. Highly specialized cases or complex emergencies will require evacuation. The U.S. is a frequent destination.

The national emergency number is 911.

Hyperbaric chambers for diving injuries are located in Guayaquil and Puerto Ayora.

Upfront payment by cash or credit card, up to the total of all anticipated charges, is generally required by hospitals catering to foreigners prior to services or treatment. Upfront payment of other than a modest deposit may be waived by hospitals that have existing cashless agreements with at least some major international insurance providers. Public hospitals may provide some services free to foreigners.

Travax content represents decision-relevant, expert synthesis of real-time data reconciled with new and existing available advice from authoritative national and international bodies. Recommendations may differ from those of individual countries' public health authorities. Travax country-specific recommendations pertain to healthy adult travelers. Guidance regarding pediatric and special needs travelers can be found under the relevant topic in the Travax Library.

Last changed April 23, 2018

© 2018 Shoreland, Inc. All rights reserved.